arXiv:1403.6061v1 [math.AG] 24 Mar 2014

Degenerations of Kählerian K3 surfaces with finite symplectic automorphism groups

Viacheslav V. Nikulin

Abstract

Using results of our preprint "Kählerian K3 surfaces and Niemeier lattices" arXiv:1109.2879 (and the corresponding papers), we classify degenerations of Kählerian K3 surfaces with finite symplectic automorphism groups.

Dedicated to the memory of Andrey Todorov

1 Introduction

Using results of our preprint "Kählerian K3 surfaces and Niemeier lattices" arXiv:1109.2879 [16] (and the corresponding papers), we classify degenerations of Kählerian K3 surfaces with finite symplectic automorphism groups.

Here we mainly consider cases which can be marked by the Niemeier lattices N_{22} and N_{23} and degenerations of codimension 1. See Section 3 for more details.

We hope to consider all cases in further variants of the paper and further publications and give more details.

This was the subject of my talk on the conference "Automorphic forms, Lie algebras and String Theory" at Lille University on 4 March 2014

2 Existence of a primitive embedding of an even lattice into even unimodular lattices, according to [14]

In this paper, we use notations, definitions and results of [14] about lattices (that is non-degenerate integral (over $\mathbb{Z}$) symmetric bilinear forms). In particular, $\oplus$ denotes the orthogonal sum of lattices, quadratic forms. For a prime p, we denote by $\mathbb{Z}_p$ the ring of p-adic integers, and by $\mathbb{Z}_p^*$ its group of invertible elements.

Let S be a lattice. Let $A_S = S^*/S$ be its discriminant group, and q_S its discriminant quadratic form on A_S where we assume that the lattice S is even: that is x^2 is even for any $x \in S$. We denote by $l(A_S)$ the minimal number of generators of the finite Abelian group A_S, and by $|A_S|$ its order. For a prime p, we denote by $q_{S_p} = q_{S \otimes \mathbb{Z}_p}$ the p-component of q_S (equivalently, the discriminant quadratic form of the p-adic lattice $S \otimes \mathbb{Z}_p$). A quadratic form on a group of order 2 is denoted by $q_\theta^{(2)}(2)$. A p-adic lattice $K(q_{S_p})$ or the rank $l(A_{S_p})$ with the discriminant quadratic form q_{S_p} is denoted by $K(q_{S_p})$. It is unique, up to isomorphisms, for $p \neq 2$, and for $p = 2$, if $q_{S_2} \ncong q_\theta^{(2)}(2) \oplus q'$. We have the following result where an embedding $S \subset L$ of lattices is called *primitive* if L/S has no torsion.

Theorem 1. *(Theorem 1.12.2 in [14]).*

Let S be an even lattice of the signature $(t_{(+)}, t_{(-)})$, and $l_{(+)}$, $l_{(-)}$ are integers.

Then, there exists a primitive embedding of S into one of even unimodular lattices of the signature $(l_{(+)}, l_{(-)})$ if and only if the following conditions satisfy:

(1) $l_{(+)} - l_{(-)} \equiv 0 \bmod 8$;

(2) $l_{(+)} - t_{(+)} \geq 0$, $l_{(-)} - t_{(-)} \geq 0$, $l_{(+)} + l_{(-)} - t_{(+)} - t_{(-)} \geq l(A_S)$;

(3) $(-1)^{l_{(+)} - t_{(+)}} |A_S| \equiv \det K(q_{S_p}) \bmod (\mathbb{Z}_p{}^*)^2$ for each odd prime p such that $l_{(+)} + l_{(-)} - t_{(+)} - t_{(-)} = l(A_{S_p})$;

(4) $|A_S| \equiv \pm \det K(q_{S_2}) \bmod (\mathbb{Z}_2{}^*)^2$, if $l_{(+)} + l_{(-)} - t_{(+)} - t_{(-)} = l(A_{S_2})$ and $q_{S_2} \ncong q_\theta^{(2)}(2) \oplus q'$.

Remark that if the last inequality in (2) is strict then one does not need the conditions (3) and (4). If $q_{S_2} \cong q_\theta^{(2)}(2) \oplus q'$, then one does not need the condition (4).

3 Example related to Kummer surfaces, and the general theory.

Let us consider Kählerian K3 surfaces which are Kählerian Kummer sufaces. They are minimal desingularizations of the quotients of 2-dimensional complex tori by the involution of the inversion. A general Kummer surface X has the Picard number 16 and has 16 non-singular rational curves which define the Dynkin diagram $16\mathbb{A}_1$. Moreover, X has the finite symplectic automorphism group $(C_2)^4$. General Kählerian K3 surfaces with such symplectic automorphism group $(C_2)^4$ have Picard number 15. See [11], [13]. Thus Kummer surfaces can be considered as the *degeneration of codimension* 1 of Kählerian K3 surfaces with symplectic automorphism group $(C_2)^4$.

This classical example raises the following interesting questions.

Do Kählerian K3 surfaces with finite symplectic automorphism group $(C_2)^4$ have other degenerations of codimension 1 which are different from Kummer?

What are degenerations of codimension 1 (and other codimensions) of K3 surfaces with other finite symplectic automorphism groups?

Using results of our preprint [16], we can answer this questions.

For example, we show that Kählerian K3 surfaces with finite symplectic automorphism group $(C_2)^4$ have another degeneration of codimension 1 which has the type $4\mathbb{A}_1$. Thus, the Picard lattice of the K3 surface has the rank 16 and the K3 surface has exactly 4 non-singular rational curves which have the Dynkin diagram $4\mathbb{A}_1$ and it has the automorphism group $(C_2)^4$. It follows from Theorem 4 below.

Using markings of Kählerian K3 surfaces by Niemeier lattices which was developed in [16], this is reduced to the following question. We use notations and results from [16].

Let N_i be one of 23 Niemeier lattices which have non-trivial root systems. We fix the bases $P(N_i)$ of the root system and consider the automorphism group $A(N_i)$ of N_i which preserves the bases $P(N_i)$. Let $H \subset A(N_i)$ be a subgroup of $A(N_i)$. We denote by $(N_i)_H = (N_i^H)^{\perp}_{N_i} \subset N_i$ the coinvariant sublattice of H. We say that H is *maximal* if $H = Clos(H)$ is the maximal subgroup of $A(N_i)$ with the same coinvariant sublattice N_H. We can restrict considering the maximal case only. A subgroup H is *Kählerian K3 automorphism group* if N_H has a primitive embedding into the lattice L_{K3} which is an even unimodular lattice of the signature $(3, 19)$. It is isomorphic to the lattice $H^2(X, \mathbb{Z})$ of Kählerian K3 surfaces. Equivalently, N_H satisfies conditions of Theorem 1 for $l_{(+)} = 3$, $l_{(-)} = 19$.

Let $\alpha \in P(N_i)$ and $H(\alpha)$ is the orbit of H for $\alpha \in P(N_i)$. The primitive sublattice $S = [N_H, \alpha]_{pr} \subset N_i$ contains $H(\alpha)$ and $H(\alpha)$ gives the basis of the system of roots with square (-2) in S. If S has a primitive embedding into L_{K3} (equivalently, if S satisfies Theorem 1 for $l_{(+)} = 3$, $l_{(-)} = 19$, then S can be considered as Picard lattice of general Kählerian K3 surfaces with symplectic automorphism group H and classes of non-singular rational curves $H(\alpha)$. Here we use the epimorphicity of the Torelli map for Kählerian K3 surfaces, see A. Todorov [21] and Y. Siu [19]. Since $\operatorname{rk} S = \operatorname{rk} N_H + 1$, these K3 surfaces can be considered as degeneration of codimension 1 of Kählerian K3 surfaces with the automorphism group H. The primitive embedding $S \subset N_i$ can be considered as *marking* of the Kählerian K3 surfaces by the Niemeier lattice N_i. See [16] for details.

In [16] we described Kählerian K3 conjugacy classes for Niemeier lattices N_i, $i = 1, \dots, 23$. Thus, using these classification and Theorem 1, we can describe degenerations of codimension 1 of Kählerian K3 surfaces with finite symplectic automorphism groups. By considering several orbits of H, we can describe all degenerations.

For example, the group $(C_2)^4$ corresponds to the following Kählerian K3 conjugacy classes for N_{23}. This case can be marked by N_{23} only. See [16]. We use the basis $\alpha_1, \dots, \alpha_{24}$ for the

roots system $24\mathbb{A}_1$ of N_{23}.

n=21, $H \cong C_2^4$ ($|H| = 16$, $i = 14$): $\operatorname{rk} N_H = 15$, $(N_H)^*/N_H \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$.

$$H_{21,1} = [(\alpha_2\alpha_{20})(\alpha_3\alpha_{10})(\alpha_5\alpha_6)(\alpha_8\alpha_{11})(\alpha_9\alpha_{21})(\alpha_{12}\alpha_{22})(\alpha_{17}\alpha_{23})(\alpha_{19}\alpha_{24}),$$

$$(\alpha_2\alpha_{19})(\alpha_3\alpha_5)(\alpha_6\alpha_{10})(\alpha_8\alpha_9)(\alpha_{11}\alpha_{21})(\alpha_{12}\alpha_{23})(\alpha_{17}\alpha_{22})(\alpha_{20}\alpha_{24}),$$

$$(\alpha_1\alpha_{16})(\alpha_2\alpha_{20})(\alpha_3\alpha_6)(\alpha_5\alpha_{10})(\alpha_{12}\alpha_{23})(\alpha_{14}\alpha_{18})(\alpha_{17}\alpha_{22})(\alpha_{19}\alpha_{24}),$$

$$(\alpha_1\alpha_{14})(\alpha_2\alpha_{24})(\alpha_3\alpha_5)(\alpha_6\alpha_{10})(\alpha_{12}\alpha_{22})(\alpha_{16}\alpha_{18})(\alpha_{17}\alpha_{23})(\alpha_{19}\alpha_{20})]$$

with orbits $\{\alpha_1, \alpha_{16}, \alpha_{14}, \alpha_{18}\}, \{\alpha_2, \alpha_{20}, \alpha_{19}, \alpha_{24}\}, \{\alpha_3, \alpha_{10}, \alpha_5, \alpha_6\}, \{\alpha_8, \alpha_{11}, \alpha_9, \alpha_{21}\}$, $\{\alpha_{12}, \alpha_{22}, \alpha_{23}, \alpha_{17}\}$;

$$H_{21,2} = [(\alpha_1\alpha_3)(\alpha_2\alpha_{23})(\alpha_5\alpha_{14})(\alpha_6\alpha_{16})(\alpha_{10}\alpha_{18})(\alpha_{12}\alpha_{20})(\alpha_{17}\alpha_{24})(\alpha_{19}\alpha_{22}),$$

$$(\alpha_1\alpha_2)(\alpha_3\alpha_{23})(\alpha_5\alpha_{17})(\alpha_6\alpha_{12})(\alpha_{10}\alpha_{22})(\alpha_{14}\alpha_{24})(\alpha_{16}\alpha_{20})(\alpha_{18}\alpha_{19}),$$

$$(\alpha_1\alpha_{16})(\alpha_2\alpha_{20})(\alpha_3\alpha_6)(\alpha_5\alpha_{10})(\alpha_{12}\alpha_{23})(\alpha_{14}\alpha_{18})(\alpha_{17}\alpha_{22})(\alpha_{19}\alpha_{24}),$$

$$(\alpha_1\alpha_{14})(\alpha_2\alpha_{24})(\alpha_3\alpha_5)(\alpha_6\alpha_{10})(\alpha_{12}\alpha_{22})(\alpha_{16}\alpha_{18})(\alpha_{17}\alpha_{23})(\alpha_{19}\alpha_{20})]$$

with orbits $\{\alpha_1, \alpha_3, \alpha_2, \alpha_{16}, \alpha_{14}, \alpha_{23}, \alpha_6, \alpha_5, \alpha_{20}, \alpha_{24}, \alpha_{18}, \alpha_{12}, \alpha_{17}, \alpha_{10}, \alpha_{19}, \alpha_{22}\}$ (we give only orbits with more than one element).

The group $H_{21,1}$ and its orbits with 4 elements give degenerations of the type $4\mathbb{A}_1$ (only for these orbits the lattice S has a primitive embedding into L_{K3} by Theorem 1). The group $H_{21,2}$ and its orbit with 16 elements gives the degeneration of the type $16\mathbb{A}_1$ which gives the case of Kummer surfaces (only for this orbit the lattice S has a primitive embedding into L_{K3} by Theorem 1). Thus Kählerian K3 surfaces with symplectic automorphism group $H \cong (C_2)^4$ have codimension 1 degenerations only of the types $4\mathbb{A}_1$ and $16\mathbb{A}_1$. We also calculate the discriminant group $A_S = S^*/S$ of S for both cases.

Similarly below we describe degenerations of codimension 1 which are marked by N_{23} (Theorem 2) and by N_{22} (Theorem 3) and degenerations of codimension 1 which can be marked by these Niemeier lattices only (Theorem 4)

We hope to consider all other cases and degenerations of arbitrary codimension in further variants of this paper and further publications.

4 Classification of degenerations of finite symplectic automorphism groups of Kählerian K3 surfaces which can be marked by the Niemeier lattice N_{23}

Here we mark by $n = 81, \dots, 1$ types of abstract finite groups of symplectic automorphisms of Kählerian K3 surfaces. See [13], [9], [22] and [6].

Theorem 2. *Types of degenerations of codimension one $N_H \subset S \subset N_{23}$ of maximal ($Clos(H) = H$) finite symplectic automorphism groups of Kählerian K3 surfaces which can be marked by the Niemeier lattice N_{23} are shown in the Table 1 below. They are given by orbits (orbits of α_i) of Kählerian K3 conjugation classes of $A(N_{23})$ for the Niemeier lattice N_{23} classified in [16]. We use notations from [16]. We give the type of the root system and the isomorphism class of the discriminant group A_S of the corresponding lattice $S = [N_H, \alpha_i]_{pr}$.*

Table 1: All possible markings by N_{23} of degenerations of codimension one of finite symplectic automorphism groups of Kählerian K3 surfaces.

n=81, $H \cong M_{20}$ ($|H| = 960$, $i = 11357$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=80, $H \cong F_{384}$ ($|H| = 384$, $i = 18135$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=79, $H \cong \mathfrak{A}_6$ ($|H| = 360$, $i = 118$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=78, $H \cong \mathfrak{A}_{4,4}$ ($|H| = 288$, $i = 1026$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=77, $H \cong T_{192}$ ($|H| = 192$, $i = 1493$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=76, $H \cong H_{192}$ ($|H| = 192$, $i = 955$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=75, $H \cong 4^2\mathfrak{A}_4$ ($|H| = 192$, $i = 1023$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong (\mathbb{Z}/8\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations:* Type $16\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/8\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{75,1}$ with the orbit of α_1).

n=74, $H \cong L_2(7)$ ($|H| = 168$, $i = 42$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=70, $H \cong \mathfrak{S}_5$ ($|H| = 120$, $i = 34$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=65, $H \cong 2^4D_6$ ($|H| = 96$, $i = 227$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations:* $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2$ ($H_{65,3}$ with the orbit of α_3 or α_{10}; $H_{65,4}$ with the orbit of α_3, α_4 or α_7); $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{65,2}$ with the obit of α_1; $H_{65,4}$ with the orbit of α_1); $12\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^3$ ($H_{65,3}$ with the orbit of α_1); $16\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{65,1}$ with the orbit of α_1).

n=63, $H \cong M_9$ ($|H| = 72$, $i = 41$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=62, $H \cong N_{72}$ ($|H| = 72$, $i = 40$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=61, $H \cong \mathfrak{A}_{4,3}$ ($|H| = 72$, $i = 43$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong (\mathbb{Z}/12\mathbb{Z})^2 \times \mathbb{Z}/3\mathbb{Z}$. *Degenerations:* $3\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/12\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{61,1}$ with the orbit of α_7 or α_{16}); $12\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ ($H_{61,1}$ with the orbit of α_2).

n=56, $H \cong \Gamma_{25}a_1$ ($|H| = 64$, $i = 138$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^3$. *Degenerations:* $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2$ ($H_{56,1}$ with the orbit of α_1 or α_6); $16\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^3$ ($H_{56,2}$ with the orbit of α_2).

n=55, $H \cong \mathfrak{A}_5$ ($|H| = 60$, $i = 5$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong \mathbb{Z}/30\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$. *Degenerations:* $\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/30\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{55,1}$ with the orbit of α_1, α_{10} or α_{24}; $H_{55,2}$ with the orbit of α_1, α_5, α_{21} or α_{24}); $5\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/30\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{55,1}$ with the orbit of α_1, α_{10} or α_{24}; $H_{55,2}$ with the orbit of α_1, α_5, α_{21} or α_{24}); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/20\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z}$ ($H_{55,1}$ with the orbit of α_4); $10\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/60\mathbb{Z}$ ($H_{55,1}$ with the orbit of α_3); $15\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/10\mathbb{Z}(\mathbb{Z}/2\mathbb{Z})^2$ ($H_{55,2}$ with the orbit of α_2).

n=54, $H \cong T_{48}$ ($|H| = 48$, $i = 29$): $\operatorname{rk} N_H = 19$ and $(N_H)^*/N_H \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. *No degenerations.*

n=51, $H \cong C_2 \times \mathfrak{S}_4$ ($|H| = 48$, $i = 48$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong (\mathbb{Z}/12\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations:* $2\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/12\mathbb{Z})^2 \times \mathbb{Z}/4\mathbb{Z}$ ($H_{51,2}$ with the orbit of α_2 or α_{10}; $H_{51,3}$ with the orbit of α_2; $H_{51,4}$ with the orbit of α_3 or α_6); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{51,2}$ with the orbit of α_1; $H_{51,4}$ with the orbit of α_1 or α_5); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2$ ($H_{51,3}$ with the orbit of α_4; $H_{51,4}$ with the orbit of α_2 or α_7); $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{51,1}$ with the orbit of α_1 or α_4; $H_{51,3}$ with the orbit of α_1); $12\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{51,2}$ with the orbit of α_4).

n=49, $H \cong 2^4C_3$ ($|H| = 48$, $i = 50$): $\operatorname{rk} N_H = 17$, $(N_H)^*/N_H \cong \mathbb{Z}/24\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^4$. *Degenerations of codimension* 1*:* $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{49,1}$ with the orbit of α_2 or α_4; $H_{49,3}$ with the orbit of α_2, α_4, α_8, α_{10} or α_{11}); $12\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{49,1}$ with the orbit of α_1); $16\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^3$ ($H_{49,2}$ with the orbit of α_2).

n=48, $H \cong \mathfrak{S}_{3,3}$ ($|H| = 36$, $i = 10$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong \mathbb{Z}/18\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^2$. *Degenerations:* $3\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/18\mathbb{Z} \times (\mathbb{Z}/6\mathbb{Z})^2$ ($H_{48,1}$ with the orbit of α_5 or α_7); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/36\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^2$ ($H_{48,1}$ with the orbit of α_2); $9\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^3$ ($H_{48,1}$ with the orbit of α_4).

n=46, $H \cong 3^2C_4$ ($|H| = 36$, $i = 9$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong \mathbb{Z}/18\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$. *Degenerations:* $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/36\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ ($H_{46,1}$ with the orbit of α_1 or α_3); $9\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{46,1}$ with the orbit of α_2);

n=40, $H \cong Q_8 * Q_8$ ($|H| = 32$, $i = 49$): $\operatorname{rk} N_H = 17$, $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^5$. *Degenerations of codimension* 1*:* $8\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4$ ($H_{40,1}$ with the orbit of α_1 or α_3).

n=39, $H \cong 2^4C_2$ ($|H| = 32$, $i = 27$): $\operatorname{rk} N_H = 17$, $(N_H)^*/N_H \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations of codimension* 1*:* $4\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4$ ($H_{39,2}$ with the orbit of α_2, α_3 or α_4); $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{39,2}$ with the orbit of α_5; $H_{39,3}$ with the orbit of α_3 or α_5); $16\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{39,1}$ with the orbit of α_2).

n=34, $H \cong \mathfrak{S}_4$ ($|H| = 24$, $i = 12$): $\operatorname{rk} N_H = 17$ and $(N_H)^*/N_H \cong (\mathbb{Z}/12\mathbb{Z})^2 \times \mathbb{Z}/4\mathbb{Z}$. *Degenerations of codimension* 1*:* $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/12\mathbb{Z})^2 \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{34,1}$ with the orbit of α_5, α_6, α_{17} or α_{21}; $H_{34,2}$ with the orbit of α_6, α_{21} or α_{24}; $H_{34,3}$ with the orbit of α_5 or α_{21}; $H_{34,4}$ with the orbit of α_{10} or α_{24}); $2\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/12\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{34,2}$ with the orbit of α_4; $H_{34,3}$ with the orbit of α_4); $3\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{34,2}$ with the orbit of α_2; $H_{34,4}$ with the orbit of α_2 or α_3); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$ ($H_{34,2}$ with the orbit of α_1; $H_{34,3}$ with the orbit of α_1 or α_{10}; $H_{34,4}$ with the orbit of α_1 or α_5); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{34,1}$ with the orbit of α_1 or α_8; $H_{34,3}$ with the orbit of α_2 or α_3); $8\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/12\mathbb{Z})^2$ ($H_{34,1}$ with the orbit of α_2; $H_{34,4}$ with the orbit of α_4); $12\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ ($H_{34,2}$ with the orbit of α_3).

n=33, $H \cong C_7 \rtimes C_3$ ($|H| = 21$, $i = 1$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong (\mathbb{Z}/7\mathbb{Z})^3$. *Degenerations:* $7\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/14\mathbb{Z} \times \mathbb{Z}/7\mathbb{Z}$ ($H_{33,1}$ with the orbit of α_1, α_3 or α_{11}).

n=32, $H \cong Hol(C_5)$ ($|H| = 20$, $i = 3$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong (\mathbb{Z}/10\mathbb{Z})^2 \times \mathbb{Z}/5\mathbb{Z}$. *Degenerations:* $2\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/20\mathbb{Z} \times (\mathbb{Z}/5\mathbb{Z})^2$ ($H_{32,1}$ with the orbit of α_{19}); $5\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/10\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{32,1}$ with the orbit of α_3 or α_4); $10\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/20\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z}$ ($H_{32,1}$ with the orbit of α_2).

n=30, $H \cong \mathfrak{A}_{3,3}$ ($|H| = 18$, $i = 4$): $\operatorname{rk} N_H = 16$ and $(N_H)^*/N_H \cong \mathbb{Z}/9\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^4$. *Degenerations of codimension* 1*:* $3\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/18\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^3$ ($H_{30,1}$ with the orbit of α_4, α_5, α_{11} or α_{15}); $9\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^3$ ($H_{30,1}$ with the orbit of α_1).

n=26, $H \cong SD_{16}$ ($|H| = 16$, $i = 8$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong (\mathbb{Z}/8\mathbb{Z})^2 \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. *Degenerations:* $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/16\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{26,1}$ with the orbit of α_1 or α_3).

n=22, $H \cong C_2 \times D_8$ ($|H| = 16$, $i = 11$): $\operatorname{rk} N_H = 16$, $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^4 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations of codimension* 1*:* $2\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^5$ ($H_{22,2}$ with the orbit of α_{12} or α_{16}; $H_{22,3}$ with the orbit of α_8, α_{11}, α_{12} or α_{16}); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^4 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{22,2}$ with the orbit of α_6 or α_8; $H_{22,3}$ with the orbit of α_1, α_2, α_4 or α_6). $8\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^3 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{22,1}$ with the orbit of α_2 or α_4; $H_{22,2}$ with the orbit of α_2).

n=21, $H \cong C_2^4$ ($|H| = 16$, $i = 14$): $\operatorname{rk} N_H = 15$, $(N_H)^*/N_H \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$. *Degenerations of codimension* 1*:* $4\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^4$ ($H_{21,1}$ with the orbit of

α_1, α_2, α_3, α_8 or α_{12}); $16\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/2\mathbb{Z})^6$ ($H_{21,2}$ with the orbit of α_1).

n=18, $H \cong D_{12}$ ($|H| = 12$, $i = 4$): $\operatorname{rk} N_H = 16$ and $(N_H)^*/N_H \cong (\mathbb{Z}/6\mathbb{Z})^4$. *Degenerations of codimension* 1*:* $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^4 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{18,1}$ with the orbit of α_1 or α_6); $2\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/6\mathbb{Z})^2 \times \mathbb{Z}/3\mathbb{Z}$ ($H_{18,1}$ with the orbit of α_{15} or α_{16}); $3\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^3 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{18,1}$ with the orbit of α_4 or α_5); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/6\mathbb{Z})^2$ ($H_{18,1}$ with the orbit of α_2 or α_8).

n=17, $H \cong \mathfrak{A}_4$ ($|H| = 12$, $i = 3$): $\operatorname{rk} N_H = 16$ and $(N_H)^*/N_H \cong (\mathbb{Z}/12\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations of codimension* 1*:* $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/12\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^3$ ($H_{17,1}$ with the orbit of α_5, α_{15}, α_{17}, α_{21} or α_{24}; $H_{17,2}$ with the orbit of α_5, α_6, α_{10} or α_{17}; $H_{17,3}$ with the orbit of α_{10} or α_{15}); $3\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^3$ ($H_{17,1}$ with the orbit of α_1; $H_{17,3}$ with the orbit of α_1 or α_6); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{17,1}$ with the orbit of α_2; $H_{17,2}$ with the orbit of α_2 or α_4; $H_{17,3}$ with the orbit of α_2, α_3, α_8 or α_{11}); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2$ ($H_{17,2}$ with the orbit of α_2 or α_4); $12\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{17,1}$ with the orbit of α_3).

n=16, $H \cong D_{10}$ ($|H| = 10$, $i = 1$): $\operatorname{rk} N_H = 16$ and $(N_H)^*/N_H \cong (\mathbb{Z}/5\mathbb{Z})^4$. *Degenerations of codimension* 1*:* $\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/10\mathbb{Z} \times (\mathbb{Z}/5\mathbb{Z})^3$ ($H_{16,1}$ with the orbit of α_1, α_6, α_{19} or α_{22}); $5\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/10\mathbb{Z} \times (\mathbb{Z}/5\mathbb{Z})^2$ ($H_{16,1}$ with the orbit of α_2, α_3, α_4 or α_5).

n=12, $H \cong Q_8$ ($|H| = 8$, $i = 4$): $\operatorname{rk} N_H = 17$, $(N_H)^*/N_H \cong (\mathbb{Z}/8\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^3$.

Degenerations of codimension 1*:* $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/16\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^3$ ($H_{12,1}$ with the orbit of α_1 or α_4).

n=10, $H \cong D_8$ ($|H| = 8$, $i = 3$): $\operatorname{rk} N_H = 15$ and $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^5$. *Degenerations of codimension* 1*:* $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^5 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{10,1}$ with the orbit of α_1, α_5, α_6 or α_{24}; $H_{10,2}$ with the orbit of α_1 or α_6); $2\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{10,1}$ with the orbit of α_{10} or α_{15}; $H_{10,2}$ with the orbit of α_2, α_{10} or α_{16}); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^3$ ($H_{10,1}$ with the orbit of α_3 or α_7; $H_{10,2}$ with the orbit of α_3, α_4, α_5 or α_7); $8\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4$ ($H_{10,1}$ with the orbit of α_2).

n=9, $H \cong C_2^3$ ($|H| = 8$, $i = 5$): $\operatorname{rk} N_H = 14$, $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^6$. *Degenerations of codimension* 1: $2\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^3 \times (\mathbb{Z}/2\mathbb{Z})^4$ ($H_{9,1}$ with the orbit of α_5 or α_6; $H_{9,3}$ with the orbit of α_6, α_8, α_{10}, α_{11}, α_{12}, α_{15} or α_{16}); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$ ($H_{9,1}$ with the orbit of α_2, α_3, α_4 or α_7; $H_{9,4}$ with the orbit of α_2 or α_4); $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$ ($H_{9,2}$ with the orbit of α_2 or α_4; $H_{9,3}$ with the orbit of α_2).

n=6, $H \cong D_6$ ($|H| = 6$, $i = 1$): $\operatorname{rk} N_H = 14$ and $(N_H)^*/N_H \cong (\mathbb{Z}/6\mathbb{Z})^2 \times (\mathbb{Z}/3\mathbb{Z})^3$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^3 \times (\mathbb{Z}/3\mathbb{Z})^2$ ($H_{6,1}$ with the orbit of α_1, α_5, α_6 or α_{24}); $2\mathbb{A}_1$ with with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^4$ ($H_{6,1}$ with the orbit of α_{16});

$3\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^3 \times \mathbb{Z}/3\mathbb{Z}$ ($H_{6,1}$ with the orbit of α_3, α_7, α_{10} or α_{11}); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^3$ ($H_{6,1}$ with the orbit of α_2).

n=4, $H \cong C_4$ ($|H| = 4$, $i = 1$): $\operatorname{rk} N_H = 14$ and $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^4 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4 \times (\mathbb{Z}/2\mathbb{Z})^3$ ($H_{4,1}$ with the orbit of α_1, α_4, α_{17} or α_{18}); $2\mathbb{A}_1$ with with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^5$ ($H_{4,1}$ with the orbit of α_8 or α_{12}); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{4,1}$ with the orbit of α_2, α_3, α_7 or α_{11});

n=3, $H \cong C_2^2$ ($|H| = 4$, $i = 2$): $\operatorname{rk} N_H = 12$ and $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^6$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^7$ ($H_{3,1}$ with the orbit of α_1, α_5, α_6, α_{10}, α_{15}, α_{17}, α_{21} or α_{24}; $H_{3,2}$ with the orbit of α_1, α_5, α_6 or α_{21}); $2\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^3 \times (\mathbb{Z}/2\mathbb{Z})^4$ ($H_{3,2}$ with the orbit of α_8, α_{10}, α_{11}, α_{12}, α_{15} or α_{16}; $H_{3,3}$ with the orbit of α_1, α_2, α_3, α_4, α_5, α_6, α_7, α_8, α_{11}, α_{12}, α_{15} or α_{16}); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$ ($H_{3,1}$ with the orbit of α_2, α_3, α_4 or α_7; $H_{3,2}$ with the orbit of α_2 or α_4).

n=2, $H \cong C_3$ ($|H| = 3$, $i = 1$): $\operatorname{rk} N_H = 12$ and $(N_H)^*/N_H \cong (\mathbb{Z}/3\mathbb{Z})^6$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^5$ ($H_{2,1}$ with the orbit of α_3, α_4, α_{14}, α_{17}, α_{21} or α_{24}); $3\mathbb{A}_1$ with with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^4$ ($H_{2,1}$ with the orbit of α_1, α_2, α_5, α_6, α_{12} or α_{15}).

n=1, $H \cong C_2$ ($|H| = 2$, $i = 1$): $\operatorname{rk} N_H = 8$ and $(N_H)^*/N_H \cong (\mathbb{Z}/2\mathbb{Z})^8$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/2\mathbb{Z})^9$ ($H_{1,1}$ with the orbit of α_1, α_4, α_8, α_{12}, α_{15}, α_{17}, α_{18} or α_{21}); $2\mathbb{A}_1$ with with $A_S \cong \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$ ($H_{1,1}$ with the orbit of α_2, α_3, α_5, α_6, α_7, α_{10}, α_{11} or α_{14}).

5 Classification of degenerations of finite symplectic automorphism groups of Kählerian K3 surfaces which can be marked by the Niemeier lattice N_{22}

Here we mark by $n = 81, \dots, 1$ types of abstract finite groups of symplectic automorphisms of Kählerian K3 surfaces. See [13], [9], [22] and [6].

Theorem 3. *Types of degenerations of codimension one $N_H \subset S \subset N_{22}$ of maximal ($Clos(H) = H$) finite symplectic automorphism groups of Kählerian K3 surfaces which can be marked by the Niemeier lattice N_{22} are shown in the Table 2 below. They are given by orbits (orbits of $\alpha_{i,j}$) of Kählerian K3 conjugation classes of $A(N_{22})$ for the Niemeier lattice N_{22} classified in [16]. We use notations from [16]. We give the type of the root system and the isomorphism class of the discriminant group A_S of the corresponding lattice $S = [N_H, \alpha_{i,j}]_{pr}$.*

Table 2: All possible markings by N_{22} of degenerations of codimension one of finite symplectic automorphism groups of Kählerian K3 surfaces.

n=79, $H \cong \mathfrak{A}_6$ ($|H| = 360$, $i = 118$): $\operatorname{rk} N_H = 19$ and $(N_H)^*/N_H \cong \mathbb{Z}/60\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$. *No degenerations.*

n=70, $H \cong \mathfrak{S}_5$ ($|H| = 120$, $i = 34$): $\operatorname{rk} N_H = 19$ and $(N_H)^*/N_H \cong \mathbb{Z}/60\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z}$. *No degenerations.*

n=63, $H \cong M_9$ ($|H| = 72$, $i = 41$): $\operatorname{rk} N_H = 19$ and $(N_H)^*/N_H \cong \mathbb{Z}/18\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. *No degenerations.*

n=62, $H \cong N_{72}$ ($|H| = 72$, $i = 40$): $\operatorname{rk} N_H = 19$, $(N_H)^*/N_H \cong \mathbb{Z}/36\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^2$. *No degenerations.*

n=55, $H \cong \mathfrak{A}_5$ ($|H| = 60$, $i = 5$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong \mathbb{Z}/30\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z}$. *Degenerations*: $\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/30\mathbb{Z} \times \mathbb{Z}/10\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{55,1}$ with the orbit of $\alpha_{1,10}$ or $\alpha_{2,10}$; $H_{55,2}$ with the orbit of $\alpha_{1,6}$, $\alpha_{2,6}$, $\alpha_{1,8}$ or $\alpha_{2,8}$); $5\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/30\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{55,1}$ with the orbit of $\alpha_{1,6}$ or $\alpha_{2,6}$); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/20\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z}$ ($H_{55,1}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$); $10\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/60\mathbb{Z}$ ($H_{55,2}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$).

n=54, $H \cong T_{48}$ ($|H| = 48$, $i = 29$): $\operatorname{rk} N_H = 19$ and $(N_H)^*/N_H \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. *No degenerations.*

n=48, $H \cong \mathfrak{S}_{3,3}$ ($|H| = 36$, $i = 10$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong \mathbb{Z}/18\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^2$. *Degenerations*: $3\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/18\mathbb{Z} \times (\mathbb{Z}/6\mathbb{Z})^2$ ($H_{48,2}$ with the orbit of $\alpha_{1,6}$, $\alpha_{2,6}$, $\alpha_{1,7}$ or $\alpha_{2,7}$); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/36\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^2$ ($H_{48,2}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{1,2}$); $9\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^3$ ($H_{48,1}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$).

n=46, $H \cong 3^2C_4$ ($|H| = 36$, $i = 9$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong \mathbb{Z}/18\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$. *Degenerations*: $9\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{46,1}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$); $9\mathbb{A}_2$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ ($H_{46,2}$ with the orbit of $\alpha_{1,1}$).

n=34, $H \cong \mathfrak{S}_4$ ($|H| = 24$, $i = 12$): $\operatorname{rk} N_H = 17$ and $(N_H)^*/N_H \cong (\mathbb{Z}/12\mathbb{Z})^2 \times \mathbb{Z}/4\mathbb{Z}$. *Degenerations of codimension* 1: $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/12\mathbb{Z}$ ($H_{34,1}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$); $6\mathbb{A}_2$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ ($H_{34,1}$ with the orbit of $\alpha_{1,2}$).

n=32, $H \cong Hol(C_5)$ ($|H| = 20$, $i = 3$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong (\mathbb{Z}/10\mathbb{Z})^2 \times \mathbb{Z}/5\mathbb{Z}$. *Degenerations*: $5\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/10\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{32,1}$ with the orbit of $\alpha_{1,3}$ or $\alpha_{2,3}$); $5\mathbb{A}_2$ with $A_S \cong \mathbb{Z}/10\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z}$ ($H_{32,1}$ with the orbit of $\alpha_{1,6}$).

n=30, $H \cong \mathfrak{A}_{3,3}$ ($|H| = 18$, $i = 4$): $\operatorname{rk} N_H = 16$ and $(N_H)^*/N_H \cong \mathbb{Z}/9\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^4$. *Degenerations of codimension* 1: $3\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/18\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^3$ ($H_{30,2}$ with the orbit of $\alpha_{1,1}$, $\alpha_{2,1}$, $\alpha_{1,2}$, $\alpha_{2,2}$, $\alpha_{1,6}$, $\alpha_{2,6}$, $\alpha_{1,7}$ or $\alpha_{2,7}$); $9\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^3$ ($H_{30,1}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$).

n=26, $H \cong SD_{16}$ ($|H| = 16$, $i = 8$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong (\mathbb{Z}/8\mathbb{Z})^2 \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. *Degenerations*: $2\mathbb{A}_2$ with $A_S \cong (\mathbb{Z}/8\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{26,1}$ with the orbit of $\alpha_{1,3}$); $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/16\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{26,1}$ with the orbit of $\alpha_{1,5}$ or $\alpha_{2,5}$).

n=18, $H \cong D_{12}$ ($|H| = 12$, $i = 4$): $\operatorname{rk} N_H = 16$ and $(N_H)^*/N_H \cong (\mathbb{Z}/6\mathbb{Z})^4$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^4 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{18,1}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$); $2\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/6\mathbb{Z})^2 \times \mathbb{Z}/3\mathbb{Z}$ ($H_{18,1}$ with the orbit of $\alpha_{1,7}$ or $\alpha_{2,7}$); $3\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^3 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{18,1}$ with the orbit of $\alpha_{1,2}$ or $\alpha_{2,2}$); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/6\mathbb{Z})^2$ ($H_{18,1}$ with the orbit of $\alpha_{1,5}$ or $\alpha_{2,5}$).

n=17, $H = \mathfrak{A}_4$ ($|H| = 12$, $i = 3$): $\operatorname{rk} N_H = 16$ and $(N_H)^*/N_H \cong (\mathbb{Z}/12\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/12\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^3$ ($H_{17,1}$ with the orbit of $\alpha_{1,1}$, $\alpha_{2,1}$, $\alpha_{1,2}$ or $\alpha_{2,2}$); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ($H_{17,1}$ with the orbit of $\alpha_{1,3}$ or $\alpha_{2,3}$); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2$ ($H_{17,1}$ with the orbit of $\alpha_{1,5}$ or $\alpha_{2,5}$).

n=16, $H \cong D_{10}$ ($|H| = 10$, $i = 1$): $\operatorname{rk} N_H = 16$ and $(N_H)^*/N_H \cong (\mathbb{Z}/5\mathbb{Z})^4$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/10\mathbb{Z} \times (\mathbb{Z}/5\mathbb{Z})^3$ ($H_{16,1}$ with the orbit of $\alpha_{1,1}$, $\alpha_{2,1}$, $\alpha_{1,2}$ or $\alpha_{2,2}$); $5\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/10\mathbb{Z} \times (\mathbb{Z}/5\mathbb{Z})^2$ ($H_{16,1}$ with the orbit of $\alpha_{1,3}$, $\alpha_{2,3}$, $\alpha_{1,5}$ or $\alpha_{2,5}$).

n=12, $H \cong Q_8$ ($|H| = 8$, $i = 4$): $\operatorname{rk} N_H = 17$, $(N_H)^*/N_H \cong (\mathbb{Z}/8\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^3$. *Degenerations of codimension* 1: $\mathbb{A}_2$ with $A_S \cong (\mathbb{Z}/8\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{12,1}$ with the orbit of $\alpha_{1,5}$, $\alpha_{1,8}$ or $\alpha_{1,11}$); $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/16\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^3$ ($H_{12,1}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$).

n=10, $H \cong D_8$ ($|H| = 8$, $i = 3$): $\operatorname{rk} N_H = 15$ and $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^5$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^5 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{10,1}$ with the orbit of $\alpha_{1,1}$, $\alpha_{2,1}$, $\alpha_{1,2}$ or $\alpha_{2,2}$); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^3$ ($H_{10,1}$ with the orbit of $\alpha_{1,3}$, $\alpha_{2,3}$, $\alpha_{1,5}$ or $\alpha_{2,5}$); $2\mathbb{A}_2$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4$ ($H_{10,1}$ with the orbit of $\alpha_{1,9}$).

n=6, $H \cong D_6$ ($|H| = 6$, $i = 1$): $\operatorname{rk} N_H = 14$, $(N_H)^*/N_H \cong (\mathbb{Z}/6\mathbb{Z})^2 \times (\mathbb{Z}/3\mathbb{Z})^3$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^3 \times (\mathbb{Z}/3\mathbb{Z})^2$ ($H_{6,1}$ with the orbit of $\alpha_{1,8}$ or $\alpha_{2,8}$; $H_{6,2}$ with the orbit of $\alpha_{1,6}$, $\alpha_{2,6}$, $\alpha_{1,8}$, $\alpha_{2,8}$, $\alpha_{1,10}$ or $\alpha_{2,10}$); $2\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^4$ ($H_{6,1}$ with the orbit of $\alpha_{1,6}$ or $\alpha_{2,6}$); $3\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^3 \times \mathbb{Z}/3\mathbb{Z}$ ($H_{6,2}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{2,1}$); $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^3$ ($H_{6,2}$ with the orbit of $\alpha_{1,2}$ or $\alpha_{2,2}$).

n=4, $H \cong C_4$ ($|H| = 4$, $i = 1$): $\operatorname{rk} N_H = 14$ and $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^4 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4 \times (\mathbb{Z}/2\mathbb{Z})^3$ ($H_{4,1}$ with the orbit of $\alpha_{1,2}$, $\alpha_{2,2}$, $\alpha_{1,5}$ or $\alpha_{2,5}$); $\mathbb{A}_2$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4 \times \mathbb{Z}/2\mathbb{Z}$ ($H_{4,1}$ with the orbit of $\alpha_{1,1}$ or $\alpha_{1,11}$); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$ ($H_{4,1}$ with the orbit of $\alpha_{1,3}$, $\alpha_{2,3}$, $\alpha_{1,6}$ or $\alpha_{2,6}$).

n=3, $H \cong C_2^2$ ($|H| = 4$, $i = 2$): $\operatorname{rk} N_H = 12$ and $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^6$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^7$ ($H_{3,1}$ with the orbit of

$\alpha_{1,1}$, $\alpha_{2,1}$, $\alpha_{1,2}$ or $\alpha_{2,2}$); $2\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^3 \times (\mathbb{Z}/2\mathbb{Z})^4$ ($H_{1,3}$, with the orbit of $\alpha_{1,3}$, $\alpha_{2,3}$, $\alpha_{1,5}$, $\alpha_{2,5}$, $\alpha_{1,7}$ or $\alpha_{2,7}$); $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$ ($H_{3,1}$ with the orbit of $\alpha_{1,6}$ or $\alpha_{2,6}$).

n=2, $H \cong C_3$ ($|H| = 3$, $i = 1$): $\operatorname{rk} N_H = 12$ and $(N_H)^*/N_H \cong (\mathbb{Z}/3\mathbb{Z})^6$.

Degenerations of codimension 1:

$\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^5$ ($H_{2,1}$ with the orbit of $\alpha_{1,6}$, $\alpha_{2,6}$, $\alpha_{1,8}$, $\alpha_{2,8}$, $\alpha_{1,10}$ or $\alpha_{2,10}$); $3\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/3\mathbb{Z})^4$ ($H_{2,1}$, with the orbit of $\alpha_{1,1}$, $\alpha_{2,1}$, $\alpha_{1,2}$, $\alpha_{2,2}$, $\alpha_{1,7}$ or $\alpha_{2,7}$).

n=1, $H \cong C_2$ ($|H| = 2$, $i = 1$): $\operatorname{rk} N_H = 8$ and $(N_H)^*/N_H \cong (\mathbb{Z}/2\mathbb{Z})^8$. *Degenerations of codimension* 1: $\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/2\mathbb{Z})^9$ ($H_{1,1}$ with the orbit of $\alpha_{1,1}$, $\alpha_{2,1}$, $\alpha_{1,2}$, $\alpha_{2,2}$, $\alpha_{1,5}$, $\alpha_{2,5}$, $\alpha_{1,11}$ or $\alpha_{2,11}$); $2\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$ ($H_{1,1}$ with the orbit of $\alpha_{1,3}$, $\alpha_{2,3}$, $\alpha_{1,6}$, $\alpha_{2,6}$, $\alpha_{1,7}$, $\alpha_{2,7}$, $\alpha_{1,9}$ or $\alpha_{2,9}$).

6 Classification of degenerations of finite symplectic automorphism groups of Kählerian K3 surfaces which can be marked by the Niemeier lattices N_{23} and N_{22} only

By results of Hashimoto [6] and our results in [16], maximal (that is $Clos(H) = H$) finite symplectic automorphism groups H for the Theorem below either have no degenerations since $\operatorname{rk} N_H = 19$ or can be marked by Niemeier lattices N_{23} or N_{22} only, and their degenerations are described in Theorems 2, 3. Thus, we obtain classification of their types of degenerations of codimension 1 and all their degenerations if $\operatorname{rk} N_H = 18$.

Theorem 4. *For the following maximal ($Clos(H) = H$) finite symplectic automorphism groups of Kahlerian K3 surfaces types of degenerations of codimension one $N_H \subset S$ (that is $\operatorname{rk} S = \operatorname{rk} N_H + 1$) are the following and only the following which are given in Table 3 below. More detailed description of these degenerations using markings by N_{23} and N_{22} are described in Theorems 2, 3.*

Table 3: Classification of all types of degenerations of codimension one of some finite symplectic automorphism groups of Kählerian K3 surfaces.

n=81, $H \cong M_{20}$ ($|H| = 960$, $i = 11357$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=80, $H \cong F_{384}$ ($|H| = 384$, $i = 18135$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=79, $H \cong \mathfrak{A}_6$ ($|H| = 360$, $i = 118$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=78, $H \cong \mathfrak{A}_{4,4}$ ($|H| = 288$, $i = 1026$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=77, $H \cong T_{192}$ ($|H| = 192$, $i = 1493$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=76, $H \cong H_{192}$ ($|H| = 192$, $i = 955$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=75, $H \cong 4^2\mathfrak{A}_4$ ($|H| = 192$, $i = 1023$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong (\mathbb{Z}/8\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations:* $16\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/8\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})^2$.

n=74, $H \cong L_2(7)$ ($|H| = 168$, $i = 42$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=70, $H \cong \mathfrak{S}_5$ ($|H| = 120$, $i = 34$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=65, $H \cong 2^4D_6$ ($|H| = 96$, $i = 227$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong \mathbb{Z}/24\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations:* $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z}\times(\mathbb{Z}/4\mathbb{Z})^2$, $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/24\mathbb{Z}\times(\mathbb{Z}/2\mathbb{Z})^2$, $12\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^3$, $16\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$.

n=63, $H \cong M_9$ ($|H| = 72$, $i = 41$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=62, $H \cong N_{72}$ ($|H| = 72$, $i = 40$): $\operatorname{rk} N_H = 19$. *No degenerations.*

n=56, $H \cong \Gamma_{25}a_1$ ($|H| = 64$, $i = 138$): $\operatorname{rk} N_H = 18$, $(N_H)^*/N_H \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^3$.
Degenerations: $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2$, $16\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^3$.

n=54, $H \cong T_{48}$ ($|H| = 48$, $i = 29$): $\operatorname{rk} N_H = 19$ and $(N_H)^*/N_H \cong \mathbb{Z}/24\mathbb{Z}\times\mathbb{Z}/8\mathbb{Z}\times\mathbb{Z}/2\mathbb{Z}$. *No degenerations.*

n=49, $H \cong 2^4C_3$ ($|H| = 48$, $i = 50$): $\operatorname{rk} N_H = 17$, $(N_H)^*/N_H \cong \mathbb{Z}/24\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^4$. *Degenerations of codimension* 1*:* $4\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/12\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$, $12\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$, $16\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^3$.

n=46, $H \cong 3^2C_4$ ($|H| = 36$, $i = 9$): $\operatorname{rk} N_H = 18$ and $(N_H)^*/N_H \cong \mathbb{Z}/18\mathbb{Z}\times\mathbb{Z}/6\mathbb{Z}\times\mathbb{Z}/3\mathbb{Z}$. *Degenerations:* $6\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/36\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$, $9\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/6\mathbb{Z})^2 \times \mathbb{Z}/2\mathbb{Z}$, $9\mathbb{A}_2$ with $A_S \cong \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$.

n=40, $H \cong Q_8 * Q_8$ ($|H| = 32$, $i = 49$): $\operatorname{rk} N_H = 17$, $(N_H)^*/N_H \cong (\mathbb{Z}/4\mathbb{Z})^5$. *Degenerations of codimension* 1*:* $8\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4$.

n=39, $H \cong 2^4C_2$ ($|H| = 32$, $i = 27$): $\operatorname{rk} N_H = 17$, $(N_H)^*/N_H \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$. *Degenerations of codimension* 1: $4\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^4$, $8\mathbb{A}_1$ with $A_S \cong \mathbb{Z}/8\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^2$, $16\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^2$.

n=21, $H \cong C_2^4$ ($|H| = 16$, $i = 14$): $\operatorname{rk} N_H = 15$, $(N_H)^*/N_H \cong \mathbb{Z}/8\mathbb{Z} \times (\mathbb{Z}/2\mathbb{Z})^6$. *Degenerations of codimension* 1*:* $4\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/4\mathbb{Z})^2 \times (\mathbb{Z}/2\mathbb{Z})^4$, $16\mathbb{A}_1$ with $A_S \cong (\mathbb{Z}/2\mathbb{Z})^6$.

References

[1] W. Barth, *K3 surfaces with nine cusps*, Preprint 1997, arXiv:alg-geom/9709031, 9 pages.

[2] N. Bourbaki, *Groupes et algèbres de Lie, Groupes de Coxeter et systèmes de Tits, Groupes engendres par des reflexions, systèmes de racines*, Hermann, Paris VI, 1968.

[3] D. Burns, M. Rapoport, *On the Torelli problem for Kählerian K-3 surfaces*, Ann. scient. Éc. Norm. Sup. 4^e ser. **8** (1975), 235–274.

[4] J.H. Conway, N.J.A. Sloane, *Sphere packings, lattices and groups*, Springer, 1988. 663 pages.

[5] The GAP Group, GAP-Groups, Algorithms and Programming, Version 4.6.5, 2013, http://www.gap-system.org

[6] K. Hashimoto, *Finite symplectic actions on the K3 lattice*, Preprint 2010, arXiv:1012.2682, 43 pages.

[7] Sh. Kondō, *Niemeier lattices, Mathieu groups, and finite groups of symplectic automorphisms of K3 surfaces (with appendix by Sh. Mukai)*, Duke Math. J. **92**, (1998), 593–603.

[8] Vic. S. Kulikov, *Degenerations of K3 surfaces and Enriques surfaces*, Izv. Akad. Nauk SSSR Ser. Mat. **41** (1977), no. 5, 1008–1042; English transl. in Math. USSR Izv. **11**, (1977) no. 5, 957–989.

[9] Sh. Mukai, *Finite groups of automorphisms of K3 surfaces and the Mathieu group*, Invent. math. **94**, (1988), 183–221.

[10] H.-V. Niemeier, *Definite quadratische Formen der Dimension* 24 *und Diskriminante* 1, J. Number Theory **5** (1973), 142–178.

[11] V.V. Nikulin, *On Kummer surfaces*, Izv. Akad. Nauk SSSR Ser. Mat. **39** (1975), no. 2, 278–293; English transl. in Math. USSR Izv. **9** (1975), 261–275.

[12] V.V. Nikulin, *Finite automorphism groups of Kählerian surfaces of type K3*, Uspehi matem. nauk **31** (1976), no. 2, 223–224. (In Russian.)

[13] V.V. Nikulin, *Finite automorphism groups of Kähler K3 surfaces*, *Trudy Mosk. Mat. ob-va* **V. 38** (1979), 75–137; English transl. in *Trans. Moscow Math. Soc.* **V. 38** (1980), 71–135.

[14] V.V. Nikulin, *Integral symmetric bilinear forms and some of their geometric applications*, Izv. Akad. Nauk SSSR Ser. Mat. **43** (1979), no. 1, 111–177; English transl. in Math. USSR Izv. **14** (1980), no. 1, 103–167.

[15] V.V. Nikulin, *Discrete reflection groups in Lobachevsky spaces and algebraic surfaces*, Proc. Int. Congr. Math. Berkeley 1986, Vol. 1, pp. 654-669.

[16] V.V. Nikulin, *Kahlerian K3 surfaces and Niemeier lattices*, Preprint 2011. arXiv:1109.2879v7, 77 pages .

[17] I.I. Pjatetskiĭ-Šapiro and I.R. Šafarevič, *A Torelli theorem for algebraic surfaces of type K3*, Izv. AN SSSR. Ser. mat., **35** (1971), no. 3, 530–572; English transl.: Math. USSR Izv. **5** (1971), no. 3, 547–588.

[18] I.R. Shafarevich (ed.), *Algebraic Surfaces*, Proc. Steklov Math. Inst. **75**, (1965), 3–215.

[19] Y. Siu, *A simple proof ot the surjectivity of the period map of K*3 *surfaces*, Manuscripta Math. **35** (1981), no. 3, 311–321.

[20] A. Taormina, K. Wendland, *The overarching finite symmetry groups of Kummer surfaces*, Preprint 2011, arXiv:1107.3834, 36 pages.

[21] A. Todorov, *Applications of the Kähler–Einstein–Calabi–Yau metric to moduli of K3 surfaces*, Invent. math. **61** (1981), no. 3, 251–265.

[22] G. Xiao, *Galois covers between K3 surfaces*, Ann. Inst. Fourier (Grenoble) **46** (1996), 73–88.

V.V. Nikulin
Deptm. of Pure Mathem. The University of Liverpool, Liverpool
L69 3BX, UK;
Steklov Mathematical Institute,
ul. Gubkina 8, Moscow 117966, GSP-1, Russia
vnikulin@liv.ac.uk vvnikulin@list.ru
Personal page: http://vnikulin.com